References

[1] Wagner,D.H. (1987) Equivalence of Euler and Lagrangian Equations of Gas Dynamics for Weak Solutions; J. Differential Equations, Vol.68 pp.118-136

[2] Harten,A. (1978) The Artificial Compression Method for Computation of Shocks and Contact Discontinuities: 3 Self-Adjusting Hybrid Schemes; Math. Comp., Vol.32 pp.363-389

[3] Harten,A. (1989) ENO Schemes with Subcell Resolution; J. Comput. Phys., Vol.83 pp.148-184

[4] Shu,C.W. and Osher,S. (1989) Efficient Implementation of Essentially Non-oscillatory Shock-Capturing Schemes 2; J. Comput. Phys., Vol.83 pp.32-78

[5] Liou,M.S. (1995) An extended Lagrangian method; J. Comput. Phys., Vol.118 pp.294-309

[6] Hirt,C.W., Amsden,A.A. and Cook,J.L. (1974) An Arbitrary Lagrangian-Eulerian Computing Method for All Flow Speeds; J. Comput. Phys., Vol.14 pp.227-253

[7] Pracht,W. (1975) Calculating Three-Dimensional Fluid Flows at All Speeds with an Eulerian-Lagrangian Computing Mesh; J. Comput. Phys., Vol.17 pp.132-159

[8] Margolin,L.G. (1997) Introduction to "An Arbitrary Lagrangian-Eulerian Computing Method for All Flow Speeds"; J. Comput. Phys., Vol.135 pp.198-202

[9] Hall,M.S. (1990) A Comparison of First and Second Order Rezoned and Lagrangian Godunov Solutions; J. Comput. Phys., Vol.90 pp.458-485

[10] Loh,C.Y. and Hui,W.H. (1990) A New Lagrangian Method for Steady Supersonic Flow Computation, Part 1: Godunov Scheme; J. Comput. Phys., Vol.89 pp.207-240

[11] Hui,W.H. and Loh,C.Y. (1992) A New Lagrangian Method for Steady Supersonic Flow Computation, Part 2: Slipline Resolution; J. Comput. Phys., Vol.103 pp.450-464

[12] Hui,W.H. and Loh,C.Y. (1992) A New Lagrangian Method for Steady Supersonic Flow Computation, Part 3: Strong Shocks; J. Comput. Phys., Vol.103 pp.465-471

[13] Hui,W.H. and Zhao,Y.C. (1992) A Generalized Lagrangian Method for solving the Euler Equations; *Nonlinear Hyperbolic Problems: Theoretical, Applied and Computational Aspects*, (Edited by A. Donato and F. Olivery), pp.336-345

[14] Lepage,C.Y. and Hui,W.H. (1995) A Shock-Adaptive Godunov Scheme Based on the Generalized Lagrangian Formulation; J. Comput. Phys., Vol.122 pp.291-299

[15] Hui,W.H. and Chu,D.L. (1996) Optimal Grid for the Steady Euler Equations; Comput. Fluid Dyn. J., Vol.4 pp.403-426

[16] Loh,C.Y. and Liou,M.S. (1994) A New Lagrangian Method for Three-Dimensional Steady Supersonic Flows; J. Comput. Phys., Vol.113 pp.224-248

[17] Hui,W.H. and He,Y. (1997) Hyperbolicity and Optimal Coordinates for the Three-Dimensional Supersonic Euler Equations; SIAM J. Appl. Math., Vol.57 pp.224-248

[18] W.H. Hui,W.H. (1995) Generalized Lagrangian Formulation of Computational Fluid Dynamics; *Computational Fluid Dynamics review*, M. Hafez and K. Oshima (Ed), Wiley, pp.382-398

[19] Hui,W.H., Li,P.Y. and Li,Z.W. (1999) A Unified Coordinate System for Solving the Two-Dimensional Euler Equations; To appear in J. Comput. Phys.

[20] Harten,A. and Hyman,J.M. (1983) Self-Adjusting Grid Methods for One-Dimensional Hyperbolic Conservation Laws; J. Comput. Phys., Vol.14 pp.227-253

[21] Lax,P.D. and Wendroff,B (1960) Systems of Conservation Laws; Comm. Pure Appl. Math., Vol.13 pp.217-237

[22] Ben-Artzi,M. and Falcovitz,J. (1984) A Second-Order Godunov-Type Scheme for Compressible Fluid Dynamics; J. Comput. Phys., Vol.55 pp.1-32

[23] Toro,E.F. (1997) *Riemann Solvers and Numerical Methods for Fluid Dynamics*, Springer-Verlag, Berlin, Heidelberg

[24] Woodward,P. and Colella,P. (1984) Numerical simulations of two-dimensional fluid flow with strong shocks; J. Comput. Phys., Vol.54 pp.115-173

[25] Liu,X.D. and Osher,S. (1998) Convex ENO high order multi-dimensional schemes without field by field decomposition or staggered grids; J. Comput. Phys., Vol.142 pp.304-330

[26] Von Neumann,J. and Richtmyer,R.D. (1950) A method for the numerical calculation of hydrodynamic shocks; J. Appl. Phys., Vol.21 pp.232-237

[27] Godunov,S.K. (1959) A Difference Method for Numerical Calculation of Discontinuous Solutions of the Equations of Hydrodynamics; Mat. Sb., Vol.47 pp.271-306