References

[1] W.H. Hui, P.Y. Li, and Z.W.Li, *A Unified Coordinate System for Solving the Two-Dimensional Euler Equations*, J. Comput. Phys., 153 (1999), pp. 596-637.

[2] A. Harten, *The Artificial Compression Method for Computation of Shocks and Contact Discontinuities: III Self-Adjusting Hybrid Schemes*, Math. Comp., 32 (1978), pp. 363-389.

[3] A. Harten, *ENO Schemes with Subcell Resolution*, J. Comput. Phys., 83 (1989), pp. 148-184.

[4] C. W. Shu and S. Osher, *Efficient Implementation of Essentially Non-oscillatory Shock-Capturing Schemes II*, J. Comput. Phys., 83 (1989), pp. 32-78.

[5] C. W. Hirt, A. A. Amsden and J. L. Cook, *An Arbitrary Lagrangian-Eulerian Computing Method for All Flow Speeds*, J. Comput. Phys., 14 (1974), pp. 227-253.

[6] W. Pracht, *Calculating Three-Dimensional Fluid Flows at All Speeds with an Eulerian-Lagrangian Computing Mesh*, J. Comput. Phys., 17 (1975), pp. 132-159.

[7] L. G. Margolin, *Introduction to "An Abritrary Lagrangian-Eulerian Computing Method for All Flow Speeds"*. J. Comput. Phys., 135 (1997), 198-202.

[8] M. S. Hall, *A Comparison of First and Second Order Rezoned and Lagrangian Godunov Solutions*, J. Comput. Phys., 90 (1990), 458-485.

[9] W. H. Hui and S. Koudriakov, *The Role of Coordinates in the Computation of Discontinuities in One-Dimensional Flow*, Comput. Fluid Dyn. J. 8 (2000), No. 4, 495-510.

[10] W. H. Hui and C. Y. Loh, *A New Lagrangian Method for Steady Supersonic Flow Computation, Part III: Strong Shocks*, J. Comput. Phys., 103 (1992), pp. 465-471.

[11] C. Y. Lepage and W. H. Hui, *A Shock-Adaptive Godunov Scheme Based on the Generalized Lagrangian an Formulation*, J. Comput. Phys., 122 (1995), 291-299.

[12] H. O. Kreiss and J. Lorenz, *Initial-Boundary Value Problems and the Navier-Stokes Equations*, Academic Press, San Diego, 1989.

[13] W. H. Hui and D. L. Chu, *Optimal Grid for the Steady Euler Equations*, Comput. Fluid Dyn. J., 4 (1996), pp. 403-426 (1996).

[14] M. G. Crandall and A. Majda, *Monotone Difference Approximations for Scalar Conservation Laws*, Math. Comp. 34 (1980), 1-21.

[15] G. Strang, *On the construction and Comparison of Difference Schemes*, SIAM J. Num. Anal., 5 (1968), 506-517.

[16] E.F. Toro, *Riemann problems and the WAF method for solving the two-dimensional shallow water equations*, Phil. Trans. R. Soc. Lond., A 338 (1992), pp.43-68.

[17] P. Glaister, *Approximate Riemann solutions of the shallow water equations*, J. of Hydr. Research, 26 (1988), pp.293-306.

[18] F. Alcrudo, P. Garcia-Navarro and J.-M. Saviron, *Flux difference splitting for 1D open channel flow equations*, Int. J. Numer. Meth. in Fluids, 14 (1992), pp.1009-1018.

[19] J.K. Dukowicz and B.J.A. Meltz, *Vorticity Errors in Multidimensional Lagrangian Codes*, J. Comput. Phys., 99 (1992), pp.115-134.

[20] J.Y. Yang, S.H. Chang and W.H. Hui, *A new Lagrangian method for steady super-critical shallow flow computation*, Comp. Meth. in Appl. Mech. Eng., 104 (1993), pp.333-355.

[21] J.Y. Yang, C.A. Hsu and W.H. Hui, *A generalized Lagrangian method for solving the steady shallow water equations*, Math. and Comp. in Simulation, 35 (1993), pp.43-61.

[22] F. Alcrudo and P. Garcia-Navarro, *A high-resolution Godunov-type scheme in finite volumes for the 2D shallow-water equations*, Int. J. Numer. Meth. in Fluids, 16 (1993), pp.489-505.

[23] M.E. Hubbard, *Multidimensional Slope Limiters for MUSCL-Type Finite Volume Schemes on Unstructured Grids*, J. Comput. Pyhys., 155 (1999), pp.54-74.

[24] K. Xu, *Unsplitting BGK-type Schemes for the Shallow Water Equations*, Int. J. Modern Pyhys. C, 10 No.4 (1999), pp.505-516.